


[Read ebook] Length Tension Testing Book 1, Lower Quadrant: A Workbook of Manual Therapy Techniques

Length Tension Testing Book 1, Lower Quadrant: A Workbook of Manual Therapy Techniques

Paolo Sanzo, Murray MacHutchon
ePub | *DOC | audiobook | ebooks | Download PDF



[Download](#)

[Read Online](#)

#357083 in Books 2015-02-11 Original language: English PDF # 1 9.00 x .48 x 6.061, .0 #File Name: 155059592X96 pages | File size: 55.Mb

Paolo Sanzo, Murray MacHutchon : Length Tension Testing Book 1, Lower Quadrant: A Workbook of Manual Therapy Techniques before purchasing it in order to gauge whether or not it would be worth my time, and all praised Length Tension Testing Book 1, Lower Quadrant: A Workbook of Manual Therapy Techniques:

0 of 0 people found the following review helpful. Excellent! By TOne of the most helpful physio books that I have. Highly recommended. Pictures are very helpful and everything is thoroughly explained!!

Length Tension Testing Book 1, Lower Quadrant provides clear and comprehensive descriptions and photos for assessing flexibility and length tension in the muscles of the lower quadrant. It includes techniques for the lumbar spine and pelvis, hip, knee, and foot and ankle. Each technique listing includes concise, standardized descriptions of the actions and positions involved, high-quality colour photos and alternative positions to accommodate patient variability and comfort. Most of these tests can be adapted into treatment techniques. This resource will help physiotherapists, kinesiologists, chiropractors, and massage therapists improve patient care, and it will be an invaluable reference for students at the college and university level. Also available: Length Tension Testing Book 2, Upper Quadrant.

Doody : 5 Stars, 97/100 This is a high quality resource with clear descriptions and excellent photos that promotes a clear understanding of muscle length tests for assessment and treatment in clinical practice. The format enhances the quality of each test and the size and binding enable easy reference during treatment. (Karin J Edwards, MSPT, Providence Health Services) Concise and thorough, these two manuals are essential resources for physiotherapists, athletic therapist, chiropractors and occupational therapists. (Orthopaedic Division) About the Author Dr. Paolo Sanzo is an assistant professor in the School of Kinesiology and the Northern Ontario School of Medicine at Lakehead University. A practicing physiotherapist at the Victoriaville Physiotherapy Centre in Thunder Bay, Ontario, Dr. Sanzo is also an examiner and instructor with the Orthopaedic Division of the Canadian Physiotherapy Association. Murray MacHutchon is a physiotherapist at Pembina Physiotherapy and Sports Injury Clinic in Winnipeg, Manitoba. He is also an examiner and instructor with the Orthopaedic Division of the Canadian Physiotherapy Association. Excerpt. Reprinted by permission. All rights reserved. Introduction Assessment The assessment of length tension in muscle involves the use of clinical reasoning and interpretation of the subjective and objective assessment findings. These findings include: - the referral pattern of pain; - positional findings on observation; - changes in active and passive range of motion; - findings on palpation; and - activation of the muscle and the flexibility of the muscle during the length tension assessment of the myofascial structures. Differentiation must also be made in the tension and barrier that are palpated to determine whether this is due to the myofascial tissue or the neuromeningeal tissue. Conclusions are then based on these combined tests and the muscle is determined to be normal, hypertonic, shortened or lengthened. The therapist may incorporate principles of neuromeningeal assessment to determine whether the tension and barrier present are due to the myofascial tissues or to the neuromeningeal structures. Excellent resources are available on the assessment of neuromeningeal tissue, and readers are advised to refer to these for further information and more details. Therapists must have an appreciation for the uniqueness of our anatomy. All tension testing described in this book may have to be slightly altered to accommodate the examiners or the patients anatomy. End feel The different sensations imparted to the hand of the therapist at the extremes of the passive range of motion is termed the end feel. The end feel caused by changes in the myofascial system will be different from some of the end feels associated with a joint restriction. Normal muscle at rest, and preferably with gravity eliminated, will feel soft. It will have the same feel as palpating raw steak or soft tofu. The length and tension will be as expected for the age of the patient. The contralateral muscle can be tested to confirm this. A normal muscle will contract with voluntary electromyographic (EMG) activity. Hypertonic muscle has increased elastic and viscoelastic stiffness in the absence of contractile activity. Palpation of the hypertonic muscle will feel similar to palpating well-done steak, and it will have decreased length on testing. Muscle spasm is an abnormal muscle contraction and is often painful. The EMG activity is not under voluntary control. This strong contraction will limit movement significantly. A muscle spasm is velocity dependent. If the muscle is lengthened or moved quickly, spasm will increase. Truly shortened muscle, or a muscle contracture, is often present post trauma and will feel gristly, tight and short on testing. The muscle contractile unit is shortened in the absence of EMG activity. When the muscle is lengthened or moved, the response is velocity independent. It does not matter if the movement is performed quickly or slowly, the response and length remains unchanged. The therapist must recognize the different sensations imparted to the hands at the end of the available passive range of motion and gently sense the point at which the range of motion stops. It is with our palpation skills that we determine that it is in fact the muscle being tested that is felt to be tense and that is providing the resistance to the passive movement. Both the hand providing stabilization and the hand moving the body part must together sense the tension in the muscle and the barrier present. Clinical reasoning Length tension test findings may be unrelated to the palpation findings found in muscle at rest. Therapists must base their conclusions on a clinical reasoning approach to rule out other problems or tissues at fault. Length tension testing should preferably be performed in supine lying or prone lying to unload the muscle and neutralize the effect of gravity. This cannot always be done, however, and thus alternative test positions are also described.