



[PDF] The ECG in Acute Myocardial Infarction and Unstable Angina: Diagnosis and Risk Stratification (Developments in Cardiovascular Medicine)

The ECG in Acute Myocardial Infarction and Unstable Angina: Diagnosis and Risk Stratification (Developments in Cardiovascular Medicine)

Hein J.J. Wellens, Anton M. Gorgels, P.A.F.M. Doevendans
DOC | *audiobook | ebooks | Download PDF | ePub



 Download

 Read Online

#4943824 in Books Springer 2002-11-30 Original language: English PDF # 1 10.75 x .36 x 8.251, 1.02 #File Name: 1402072147132 pages | File size: 36.Mb

Hein J.J. Wellens, Anton M. Gorgels, P.A.F.M. Doevendans : The ECG in Acute Myocardial Infarction and Unstable Angina: Diagnosis and Risk Stratification (Developments in Cardiovascular Medicine) before purchasing it in order to gauge whether or not it would be worth my time, and all praised The ECG in Acute Myocardial Infarction and Unstable Angina: Diagnosis and Risk Stratification (Developments in Cardiovascular Medicine):

0 of 0 people found the following review helpful. Four Stars By dr sanjay tiwari excellent 1 of 1 people found the

following review helpful. excellentBy Daniel BANINAthis is a must to have book.Unfortunatelly this book is from 2002.Today (2007) I'll recomend to buy other book from Prof Wellens (ECG in emergency decision making). Which is more up to date, with some little clues you will not have in this book. EG: RBBB in proximal LAD infarction.

The electrocardiogram (ECG) remains the most accessible and inexpensive diagnostic tool to evaluate the patient presenting with symptoms suggestive of acute myocardial ischemia. It plays a crucial role in decision making about the aggressiveness of therapy especially in relation to reperfusion therapy, because such therapy has resulted in a considerable reduction in mortality from acute myocardial infarction. Several factors play a role in the amount of myocardial tissue that can be salvaged by reperfusion therapy, such as the time interval between onset of coronary occlusion and reperfusion, site and size of the jeopardized area, type of reperfusion attempt (thrombolytic agent or an intracoronary catheter intervention), presence or absence of risk factors for thrombolytic agents, etc. Most important in decision making on reperfusion therapy and the type of intervention is to look for markers indicating a higher mortality rate from myocardial infarction. The ECG is a reliable, inexpensive, non-invasive instrument to obtain that information. Recently it has become clear that both in anterior and inferior myocardial infarction, the ECG frequently allows not only to identify the infarct related coronary artery, but also the site of occlusion in that artery and therefore the size of the jeopardized area. Obviously, the more proximal the occlusion, the larger the area at risk and the more aggressive the reperfusion attempt.